

<u>DB Name</u>	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u>
USPT	(pantoea) AND (fish)	7	L16
USPT	(immunity) AND (fish)[AB]	46	L15
USPT	(immunity in fish) AND (11/27/2000)	0	L14
USPT	(crustaceans and fishes feed?)[AB] AND (11/27/2000)	0	L13
USPT	(feedstuff additive)[AB] AND @AD < 11/27/2000	0	L12
USPT	(feedstuff additive)[AB] AND @AD < 11/27/2000	0	L11
USPT	('l2 (20-50'))[PN]	0	L10
USPT	('FR002608156A1' 'GB002097676A' '6190686' '6207192' '6197334' '6197288' '6190692' '6200601' '6194000' '6197329' '6200594' '6197781' '6187343' '6177122' 'D437407' '6165495' '6165511' '6173851' '6165482' '6165473' '6156738' '6153220'))[PN]	20	L9
TDBD	d l2 30-50, ti	8613	L8
TDBD	424/441	0	L7
DWPI	424/441	0	L6
EPAB	424/441	2	L5
JPAB	424/441	0	L4
PGPB	424/441	0	L3
USPT	424/441	487	L2
USPT	((424,441/)!.CCLS.)	0	L1

WEST[Help](#)[Logout](#)[Interrupt](#)[Main Menu](#)[Search Form](#)[Posting Counts](#)[Show S Numbers](#)[Edit S Numbers](#)[Preferences](#)**Search Results -**

Terms	Documents
(pantoea) AND (fish)	7

Database:

US Patents Full-Text Database
US Pre-Grant Publication Full-Text Database
JPO Abstracts Database
EPO Abstracts Database
Derwent World Patents Index
IBM Technical Disclosure Bulletins

Refine Search:

(pantoea) AND (fish)

[Clear](#)**Search History****Today's Date: 4/2/2001**

<u>DB Name</u>	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u>
USPT	('l2 (20-50)')[PN]	0	<u>L10</u>
USPT	('FR002608156A1 'GB002097676A '6190686 '6207192 '6197334 '6197288 '6190692 '6200601 '6194000 '6197329 '6200594 '6197781 '6187343 '6177122 'D437407 '6165495 '6165511 '6173851 '6165482 '6165473 '6156738 '6153220')[PN]	20	<u>L9</u>
TDBD	d l2 30-50, ti	8613	<u>L8</u>
TDBD	424/441	0	<u>L7</u>
DWPI	424/441	0	<u>L6</u>
EPAB	424/441	2	<u>L5</u>
JPAB	424/441	0	<u>L4</u>
PGPB	424/441	0	<u>L3</u>
USPT	424/441	487	<u>L2</u>
USPT	((424,441/)!.CCLS.)	0	<u>L1</u>

WEST

Structured Search

Database:

US Patents Full-Text Database	▲
US Pre-Grant Publication Full-Text Database	
JPO Abstracts Database	
EPO Abstracts Database	
Derwent World Patents Index	
IBM Technical Disclosure Bulletins	▼

Searchable Index:

<none> - All Fields	▼	Not Numeric	▼
---------------------	---	-------------	---

Term 1 text:

.immunity

Operator:

AND	▼	Proximity Distance:	1
-----	---	---------------------	---

Searchable Index:

<none> - All Fields	▼	Not Numeric	▼
---------------------	---	-------------	---

Term 2 text:

fish

Display:

10	Documents in	Display Format:	CIT	Starting With #:	1
----	--------------	-----------------	-----	------------------	---

Generate:

<input checked="" type="radio"/> Hit List	<input type="radio"/> Hit Count	<input type="radio"/> Image
-------------------------------------------	---------------------------------	-----------------------------

Search	Clear	Interrupt	Help	Logout
--------	-------	-----------	------	--------

Main Menu	Show S Numbers	Edit S Numbers	Preferences
-----------	----------------	----------------	-------------

Search History

Today's Date: 4/2/2001

Generate Collection

L15: Entry 23 of 46

File: USPT

Dec 7, 1993

US-PAT-NO: 5268357

DOCUMENT-IDENTIFIER: US 5268357 A

TITLE: Feed and feed additive for livestock, poultry and nursery fishes
for increasing disease, pseudorabies resistance

DATE-ISSUED: December 7, 1993

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Yabiki; Terutake	Sakura	N/A	N/A	JPX
Narahashi; Tsunao	Kashiwa	N/A	N/A	JPX
Okamoto; Toshihiro	Urayasu	N/A	N/A	JPX
Matsuo; Ken	Tsukuba	N/A	N/A	JPX

ASSIGNEE-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY	TYPE	CODE
National Federation of Agricultural Co-operative Associations	Tokyo	N/A	N/A	JPX		03

APPL-NO: 7/ 745846

DATE FILED: August 16, 1991

FOREIGN-APPL-PRIORITY-DATA:

COUNTRY	APPL-NO	APPL-DATE
JP	3-74841	March 13, 1991

INT-CL: [5] A23K 1/00

US-CL-ISSUED: 514/8

US-CL-CURRENT: 514/8

FIELD-OF-SEARCH: 514/8

PRIOR-ART-DISCLOSED:

U.S. PATENT DOCUMENTS

Search Selected		Search ALL	
PAT-NO	ISSUE-DATE	PATENTEE-NAME	US-CL
4079133	March 1978	Dress et al.	514/170
4777042	October 1988	Toda et al.	424/79

WEST☐ Generate Collection

L15: Entry 15 of 46

File: USPT

Jul 1, 1997

US-PAT-NO: 5643571

DOCUMENT-IDENTIFIER: US 5643571 A

TITLE: Method for controlling infectious diseases in fish and
other aquatic lifeforms in a closed culture system

DATE-ISSUED: July 1, 1997

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Sin; Y. M.	Singapore	N/A	N/A	SGX
Ling; K. H.	Singapore	N/A	N/A	SGX
Lam; T. J.	Singapore	N/A	N/A	SGX

ASSIGNEE-INFORMATION:

NAME	CITY	STATE	ZIP	CODE	COUNTRY	TYPE	CODE
National University of Singapore	N/A	N/A	N/A		SGX		03

APPL-NO: 8/ 466025

DATE FILED: June 6, 1995

PARENT-CASE:

This application is a continuation of application Ser. No.
08/103,054 filed Aug. 9, 1993, now abandoned which is a
continuation-in-part of of application Ser. No. 07/929,865 filed
Aug. 17, 1992 now abandoned.

INT-CL: [6] A61K 39/40, A61K 39/42, C07K 16/08, C07K 16/12

US-CL-ISSUED: 424/159.1; 424/164.1, 530/390.1

US-CL-CURRENT: 424/159.1; 424/164.1, 530/390.1

FIELD-OF-SEARCH: 424/130.1, 424/140.1, 424/184.1, 424/159.1,
424/164.1, 424/163.1, 530/387.1, 530/389.1, 530/389.4, 530/390.1,
530/389.5

PRIOR-ART-DISCLOSED:

U.S. PATENT DOCUMENTS

Search Selected

Search ALL

PAT-NO	ISSUE-DATE	PATENTEE-NAME	US-CL
<input type="checkbox"/> 3492400	January 1970	Klontz et al.	424/92
<input type="checkbox"/> 4309416	January 1982	Gratzek et al.	424/88
<input type="checkbox"/> 5064645	November 1991	Frechette et al.	424/85.8

OTHER PUBLICATIONS

Sin, Y.M., Ling, K.H., Lam, T.J., 1992. Protection against velvet disease in goldfish recovered from ichthyophthiriasis. *Aquaculture*, 102; 187-191.

Ling, K.H., Sin, Y.M. and Lam, T.J., 1992. Studies on immune response in freshwater ornamental fish against ichthyophthirius multifiliis fouquet 1876. *Singapore Journal of Primary Production*, 20, 46-52.

Cross "A Review of Methods to Control Ichthyophthiriasis" *Prog. Fish-culturist*, 34, 1972, pp. 165-169.

Di Conza "Relationship of Catfish Serum Antibodies to Immunoglobulin in Mucus Secretions" *Aust. Exp. Biol Med. Sci.* 49, 1971, pp. 517-519.

Dickerson "tetrahymena Pyriformis as a Protective Antigen Against Ichthyophthirius Multifiliis Infection: Comparisons Between Isolates and Ciliary Preparations" *J. Fish Biol.* 24, 1984, pp. 523-528.

Ellis "Fish Vaccination" Academic Press, London 1988 pp. 1-19.

Gratzek "control and therapy of fish diseases" *Adv. Vet. Sci. Comparat. Med.* 27, 1983, pp. 297-324.

Goven "Protection of channel Catfish, *Ictalurus Punctatus* Refinesque Against Ichthyophthirius Multifiliis Fouquet by Immunization" *J. Fish Biol.* 17, 1980, pp. 311-316.

Hines "Ichthyophthirius Multifiliis in the Mirror Carp, *Cyprinus Carpio L*" *J. Fish Biol.* 5, 1973, pp. 385-392.

Hines "Ichthyophthiriasis in the Mirror Carp *Cyprinus Carpio L*" *J. Fish Biol* 6, 1974, pp. 189-196.

Ingram "Substances Involved in the Natural Resistance of Fish to Infection -A Review" *J. Fish Biol.* 16, 1980, pp. 23-60.

Ling et al "A New Approach to Controlling Ichthyophthiriasis in a Closed Culture System of Freshwater Ornamental Fish" *Journal of Fish Diseases*, 14, 1991, pp. 595-598.

Ourth "Secretory IGM, Lysozyme and Lymphocytes in the Skin Mucus of the Channel Catfish, *Ictalurus Punctatus*" *Developmental and Comparative Immunology* vol. 4, pp. 65-74, 1980.

Post "Textbook of fish Health " Revised and Expanded T.F.H. Publications, Inc. USA, 1987 pp. 159-223.

Sakai "Opsonization by Fish Antibody and Complement in the Immune Phagocytosis by Peritoneal Exudate Cells Isolated from Salmonid Fishes" *Journal of Fish Diseases* 7, 1984, pp. 29-38.

Sin et al "Protection Against Velvet Disease in Goldfish Recovered from Ichthyophthiriasis" *Aquaculture*, 102, 1992 pp. 187-191.

Ventura et al "Histopathology of Ichthyophthirius Multifiliis Infections in Fishes" *Journal of Fish Biol.* 27, 1985, pp. 185-203.

ART-UNIT: 183

PRIMARY-EXAMINER: Nucker; Christine M.

ASSISTANT-EXAMINER: Reeves; Julie E.

ATTY-AGENT-FIRM: Lowe, Price, LeBlanc & Becker

ABSTRACT:

A method for controlling infectious diseases in fish and other aquatic lifeforms, regardless of their developmental stage. The method of controlling pathogenic organisms detrimental to aquatic lifeforms includes steps of adding fish which are pre-immunized against one or more organisms pathogenic to fish to the same aquatic medium and with normal fish. The pre-immunized fish continuously and increasingly release antibodies against the pathogenic organisms into the same aquatic medium and can protect or prevent naive (normal, non-infected) fish or naive aquatic lifeforms from infection and are able to enhance the recovery of infected fish or any aquatic lifeforms infected with the same diseases.

18 Claims, 2 Drawing figures

Exemplary Claim Number: 1

Number of Drawing Sheets: 1

BRIEF SUMMARY:

TECHNICAL FIELD

Infectious diseases cause heavy economic loss in intensive fish culture (Hines, R. S. et al., "Ichthyophthiriasis in the mirror carp *Cyprinus carpio*", III. Pathology J. Fish Dis. 6, (1973) 189-196). These may be due to the pathogenic parasites, bacteria, viruses or fungi. In Singapore, for example, the most important infectious disease in ornamental fish is due to *Ichthyophthirius multifiliis*. The protozoans (*I. multifiliis*) attack gills and skin and disturb the primary organs of excretion and osmoregulation in fish (Hines, R. S. et al., "Ichthyophthiriasis in the mirror carp *Cyprinus carpio*", III. Pathology J. Fish Dis. 6, (1973) 189-196; Ventura, M. T. et al., "Histopathology of *Ichthyophthirius multifiliis* infection in fishes", J. Fish Dis., 27, (1985) 185-203). Various methods including physical-therapy and chemotherapy have been recommended for treating *I. multifiliis* infected fish when they are showing clinical signs of the disease (Cross, D. G., "A review of methods to control *Ichthyophthiriasis*", Prog. Fish-culturist, 34, (1972) 165-169; Gratzek, J. B., "Control and therapy of fish diseases", Advances Vet. Sci. Comparat. Med., 27, (1983), 297-324; Post G. Textbook of Fish Health. (Revised and Expanded Edition), T. F. H. Publications Inc., USA, (1987)). However, all these methods can induce severe stress to the infected fish, thus rendering them more susceptible to infectious diseases subsequently. In addition, these methods are ineffective against the feeding stages of *I. multifiliis* which are associated with fish (Dickerson, H. W. et al., "Tetrahymena pyriformia as a protective

antigen against *Ichthyophthirius multifiliis* infection: comparison between isolates and ciliary preparations", J. Fish Biol., 24, (1984) 523-528). On the other hand, protection of channel catfish (Goven, B. A., et al., "Protection of channel catfish, *Ictalurus punctatus* Rafinesque, against *Ichthyophthirius multifiliis* Fouquet by immunization", J. Fish Biol., 17, (1980) 311-316; Gratzek et al., U.S. Pat. No. 4,309,416, issued Jan. 5, 1982; Dickerson, H. W. et al., "Tetrahymena pyriformis as a protective antigen against *Ichthyophthirius multifiliis* infection: comparison between isolates and ciliary preparations", J. Fish Biol., 24, (1984) 523-528) against *I. multifiliis* by immunization has been shown. Also effective protection by vaccination against bacteria (Klontz, U.S. Pat. No. 3,492,400 issued Jan. 2, 1970) and viruses (Gratzek, J. B., "Control and therapy of fish diseases", Advances Vet. Sci. Comparat. Med., 27, (1983), 297-324) have been reported in food fish. However, existing methods of administration of various vaccines are not fully practical on a mass scale. Methods of passive immunization involving injections of antibodies obtained from immunized individuals of a different species (Frechette et al., U.S. Pat. No. 5,064,645, issued Nov. 12, 1991) are also not fully practical on a large scale.

The present invention is a highly practical method wherein fish cultured in a closed recirculating system are continuously being protected against pathogens, such as protozoan parasites (e.g. *Ichthyophthirius*, *Oodinium*, *Ichthyobodo*, *Trichodina*, *Chilodonella*) and bacteria (e.g. *Aeromonas*) by one or more fish which are immunized against those pathogens and kept in the same recirculating system; these immunized fish release antibodies continuously to remove, or suppress the population of, the pathogens in the water and on/in the cultured lifeforms (if infected), and/or otherwise kill the pathogenic organisms.

SUMMARY OF THE INVENTION

A novel method of controlling pathogenic organisms in the aquatic environment and in fish and other aquatic lifeforms living in that environment, through placing in the same environment hardy fish immunized against those pathogenic organisms; these immunized fish continuously and increasingly release into the water antibodies against the pathogenic organisms concerned as long as the pathogenic organisms are still present in the water, thereby controlling the pathogenic organisms in the water and protecting the aquatic lifeforms concerned living in the same water against infection by the pathogenic organisms. In accordance with the present invention, fish may be immunized against one or more pathogenic organisms.

In the preferred embodiment for controlling pathogenic organisms in the aquatic environment, a closed culture system consisting of two (FIG. 1) or multi-tanks (FIG. 2) was used. All the tanks of this system are connected with PVC pipes with a pump and a filter so that water is continuously recirculating between the tanks. Hardy fish such as freshwater catfish or shubankin which were immunized against the pathogenic organisms such as *Ichthyophthirius* (causing white-spot disease), *Oodinium* (causing

velvet disease), Ichthyobodo, Trichodina and Chilodonella (causing slimy skin) or Aeromonas (causing fin- and gill-rot disease) were kept in one tank while the remaining tanks were used to keep naive fish or fish infected by the pathogenic organisms concerned.

The method has demonstrated its effectiveness not only in maintaining naive fish free from infection but also in facilitating the rapid recovery of the infected fish. The protection was due to the antibodies in the aquatic environment released by the immunized hardy fish.

Although evidence for the effectiveness of the invention is provided only for the white-spot disease (due to the parasitic protozoan, Ichthyophthirius), velvet disease (due to Oodinium) slimy skin disease (caused by Ichthyobodo, Trichodina and Chilodonella), and the fin- and gill-rot disease (due to the bacteria, Aeromonas) in freshwater ornamental fish, the inventive concept is applicable to any disease (whether due to parasites, bacteria, viruses, or fungi) in any species of freshwater or marine life (whether finfish or shellfish, e.g. crustaceans, mollusks and echinoderms) at any developmental stage (whether eggs, larvae, fry, juveniles or adults).

DRAWING DESCRIPTION:

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a side-view of a simple closed recirculating culture system. It consists of double deck aquarium tanks connected by PVC pipes with a pump and a filter. Water is continuously recirculating between the two tanks. The naive or infected fish (or other aquatic lifeforms) are kept in the upper tank while the immunized fish are placed in the lower tank. The lower tank is placed after the filter so that the water from the tank is pumped directing to the upper tank.

FIG. 2 is a side-view of a closed culture system which consists of more than two tanks. The main design is similar to that of FIG. 1. The lower tank D is used to keep immunized fish while the upper tanks (A to C) are used to keep naive or infected fish (or other aquatic species). Tanks A to C may be used to house naive or infected fish of different species separately.

DETAILED DESCRIPTION:

DETAILED DESCRIPTION OF THE INVENTION

The inventive method described below provides a convenient, non-toxic, reliable and effective system of disease control in an aquatic environment by using immunized fish to release antibodies to control pathogenic organisms in the aquatic environment and in fish and other aquatic lifeforms (whether freshwater or marine) in a closed culture system.

In a preferred embodiment, various species of freshwater naive or infected ornamental fish have been repeatedly tested in the closed culture system as described above (FIG. 1 and 2) with freshwater hardy fish such as catfish and shubankin which were immunized against various types of pathogenic protozoans (causing white-spot disease, velvet disease, slimy skin disease) or bacteria (causing fin- and gill-rot disease) that commonly infect ornamental fish. Under such conditions, naive fish were protected from contracting the above diseases while the already infected fish showed rapid recovery from infection. The following examples demonstrate the efficacy of the inventive method.

Materials and Methods

Naive Fish:

Four species of ornamental fish were used: angel fish (*Pterophyllum altum*), weighing 2.5-3.5 g; black goldfish (*carassius auratus*), weighing 2.5-4.5 g; platy (*Xiphophorus maculatus*), weighing 1.5-2.2 g and tiger barb (*Capoeta tetrazona*), weighing 2.0-3.5 g. In addition, hardy naive fish such as catfish (*Clarias batrachus*) weighing 150-200 g and Shubankin (*Carassius auratus*) weighing 70-100 g, were used. Unless otherwise stated, all these fish were bred from healthy broodstocks and were not previously infected with *I. multifiliis* or any other obvious parasites. The fish were kept in well-aerated aged tap water at 28.degree.-30.degree. C., pH 6.5-7.2, dissolved oxygen between 6.0-7.8 ppm, and total hardness of water (CaCO_3) 72.5-80.3 mg/l. These fish were referred to as "naive or unprimed fish". All fish were fed commercial goldfish pellets and dried tubifex worms at 1% body weight per day.

Ichthyophthirius Multifiliis:

The parasites were originally isolated from goldfish from a local fish farm and maintained by regular introduction of naive fish into the tank. "Infected fish" used for experiments were obtained by exposing 5 naive ornamental fish to infectious free-swimming tomites (1000-1500 per fish) in a 1000 ml beaker for 1 hour before transferring to the tanks of the closed culture aquarium system.

Fish Immunized Against Protozoan Parasites (Example 1 to 4 and 6):

Catfish and Shubankin were injected intraperitoneally with 0.1 ml 0.85% NaCl solution containing 2.3.times.10⁶ live tomites of *I. multifiliis*. For obtaining secondary immune response, a similar dose of live tomites was given via the same route 30 days after the initial injection. They were referred to as immunized fish.

Fish Immunized Against Bacteria (Example 5):

Catfish (*Clarias batrachus*) and Shubankin (*Carassius auratus*) were immunized by intraperitoneal injection of 0.2 ml 0.85% NaCl

solution containing 3.2.times.10.sup.6 formalin-killed *Aeromonas* sp which were collected from heavily infected tilapia and goldfish on day 0 and day 14. They were referred to as immunized fish.

Naive goldfish were infected 3 days after the exposure to the live *Aeromonas* sp. at a concentration of 3000/fish. Fish which were showing the clinical sign of the bacterial disease (fin and gill rot) were placed in the closed aquarium system.

Experiments:

All experiments were carried out in 20l glass tanks with constant aeration. No water renewal was carried out during the experimental period of 3 weeks to ensure that the fish were kept in an undisturbed environment. Fish were fed commercial goldfish pellets once a day. Unless otherwise stated, the immunized and infected fish were kept in separate tanks; the upper tank was used to keep naive or infected ornamental fish while the lower tank was used for keeping the immunized fish (FIG. 1). The number of pre-immunized, naive and infected fish per tank was varied according to the experimental design as stated in each table of results. FIG. 2 showed the same closed culture system but the number of upper tanks was increased to 3 for keeping more naive or infected ornamental fish.

Determination of Anti-multifiliis Antibodies:

Blood samples were collected by heparinized syringes and needles from the caudal vein of naive and immunized fish immediately before they were transferred into the separate tanks of the closed culture system. All sera were inactivated at 47.degree. C. for 30 minutes (Sakai, D. K., "Opsonization by fish antibody and complement in the immune phagocytosis by peritoneal exudate cells isolated from salmonid fish", *J. Fish Dis.*, 7, (1984) 29-38) before being assayed by the modified method of immobilization as suggested by Hines et al. (Hines, R. S. et al., "Ichthyophthiriasis in the mirror carp *Cyprinus carpio*", IV. Physiological dysfunction, *J. Fish Dis.*, 6, (1974) 365-371. Briefly, each serum was serially two-fold diluted with physiological saline in the cups of microtitrater plexiglass plates. About 10 to 15 active free-swimming tomites of *I. multifiliis* were introduced into each cup and the plates were gently shaken before being left in an air-conditioned room (23.degree. C.) for 48 hours. The readings of the antibody activity that could immobilize *I. multifiliis* were carried out under the light microscope. The final endpoint for the antibody activity in each diluted serum was taken when 50% of the free swimming tomites became immobilized.

The antibody activity in mucus and aquarium water was also similarly determined. Fish mucus was obtained by swabbing 10 fish individually with a soft fine sponge into 100 ml of physiological saline. It was then filtered by a fine cheese cloth to remove fish scales and other solid particles. Levels of water antibody titres were measured 4 days after both immunized and infected fish had been introduced into the double-deck aquarium system. The number of free swimming tomites that became immobilized by

the undiluted tank water was also determined.

As for Example 6, since the fish fry were far too small for collecting of blood samples for antibody titration, a modified method of immobilization testing on fish body fluid against the parasite was developed. This was done by random sampling of 3 fish fry from each of the system tank before the introduction of infective tomites, and 1 and 3 wks after the introduction of infective tomites. The fish fry were killed immediately and homogenized in a homogenizer according to their group, with 0.5 ml distilled water at 4.degree. C. The homogenized body fluid was then separated by a micro-centrifuge at 4.degree. C. The supernatant was inactivated at 47.degree. C. for 30 minutes (Sakai, D. K., "Opsonization by fish antibody and complement in the immune phagocytosis by peritoneal exudate cells isolated from salmonid fish", J. Fish Dis., 7, (1984) 29-38) before being assayed by the modified method of immobilization as described above.

Statistical analysis:

All the data were analyzed using one-way analysis of variance followed by a Student-Newman-Keuls test. A value of $p < 0.05$ was considered to be significant.

Results:

In fish farms particularly quarantine and export centers where fish are stocked for export, the qualities of fish vary considerably because fish are obtained from different suppliers. Some fish may look healthy but may be subclinically infested with a low number of infectious parasites on their body and gill surface. Because of the netting, grading, transportation, water quality and crowding, these fish were usually vulnerable to diseases. It has been well established stress caused by the above factors can lower resistance of fish to infectious diseases (Gratzek, J. B., "Control and therapy of fish diseases", Advances Vet. Sci. Comparat. Med., 27, (1983), 297-324). Although chemicals such as copper sulphate is widely used to treat ectoparasites on ornamental fish, this is not an ideal method because some species of fish are very sensitive to chemical treatment, rendering them more vulnerable to diseases subsequently. In Japan, no chemicals, except antibiotics, are allowed to treat infected culture food fish. In the U.S.A., a number of chemicals and drugs are also banned from use in aquaculture. Therefore, in order to cure infected fish and to control the outbreaks of infectious diseases, a more ideal method is to provide the aquatic environment with protective substances released from immunized fish continuously that can prevent the rapid multiplication of the disease organisms when the fish become weaker due to various stress factors.

We have shown that a significant number of infected fish recovered from infection in the presence of immunized fish (in the same tank) as compared to the controls, irrespective of species. Similar findings were also observed in the simple closed recirculating system (Example 1: Tables 1 and 2; FIGS. 1 and 2)

where immunized fish were kept in a separate tank.

EXAMPLE 1

Tables 1 and 2 show the results obtained from the closed culture system which consisted of two aquarium tanks (FIG. 1). The upper tank contained 10 infected goldfish while the lower tank contained 1 immunized catfish (Table 1) or 2 pre-immunized shubankin (Table 2). The data in both tables clearly show that immunized fish provided a significant protection for the infected fish as compared to the controls. The data further show that immunized fish after secondary immune response provided a better protection of the infected fish than those after primary immune response.

TABLE 1 Control of Ich on 10 infected goldfish in the upper tank (FIG. 1) by using a catfish, which was pre-injected with 1 (primary immunization) or 2 (secondary immunization) doses of *I. multifiliis*, in the lower tank. No. of Group recovered.^{sup.1} Survival (fish in lower No. infected rate tank) n (upper tank) (%)

Control	9	25/90	28.89	(no fish)
Control	3	7/30	23.33	(naive catfish)
Test	3	21/30	70.00	^{sup.a} (immune catfish, 5 wk after 1st injection)
Test	3	26/30	86.67	^{sup.b} (immune catfish 1 wk after 2nd injection)
Test	3	27/30	90.00	^{sup.c} (immune fish, 2 wk after 2nd injection)

n = number of replicates.

^{sup.a} $p < 0.05$, ^{sup.b} $p < 0.005$ and ^{sup.c} $p < 0.001$ significantly different from both controls. Different superscripts indicate significant difference ($p < 0.05$) between test groups. ^{sup.1} Total of the replicates.

TABLE 2 Control of Ich on 10 infected goldfish in the upper tank (FIG. 1) by using 2 shubankin, which were pre-injected with 1 (primary immunization) or 2 (secondary immunization) doses of *I. multifiliis*, in the lower tank. No. of Group recovered.^{sup.1} Survival (fish in lower No. infected rate tank) n (upper tank) (%)

Control	12	33/120	27.50	(no fish)
Control	6	16/60	26.67	(naive shubankin)
Test	6	41/60	68.33	^{sup.a} (immune shubankin, 5 wk after 1st injection)
Test	3	25/30	76.67	^{sup.b} (immune shubankin 1 wk after 2nd injection)
Test	3	28/30	86.67	^{sup.c} (immune shubankin, 2 wk after 2nd injection)

n = number of replicates.

^{sup.a} $p < 0.05$, ^{sup.b} $p < 0.005$ and ^{sup.c} $p < 0.001$ significantly different from both controls. Different superscripts indicate significant difference ($p < 0.05$) between test groups. ^{sup.1} Total of the replicates.

EXAMPLE 2

It was shown that the higher the number of immunized fish (fish which had recovered from "ich" infection or after the pathogen injection) in relation to infected fish cultured in the same tank, the better was the recovery rate of the infected fish. (Ling, Sin and Lam, September 1991, Journal of Fish Diseases, Vol 14, p. 595-598; and Ling, Sin and Lam, March 1992, Aquaculture,

Vol 102, p. 187-191, incorporated herein by reference). This indicates that the magnitude of response of the immunized fish against the parasites plays an important part in the protection of the infected fish. This view was further supported by the findings of Tables 3 and 4 that higher titres of anti-I. multifiliis antibodies were obtained in immunized fish after secondary immune response than those after primary immune response. Anti-I. multifiliis activity was also detected in the aquarium water 4 days after secondary-immunized fish were introduced into the lower tank. Although the activity in the tank water containing primary-immunized fish was only detected on day 12 based on 50% parasite immobilization test, it was detected on day 4 when the number of tomites immobilized by the tank water was counted instead of basing on 50% immobilization (Tables 5 and 6). This activity remained the same when the tank water was heated to 47.degree. C. for 30 min, indicating they are not blood complements. However, the activity was lost when the tank water was heated to 100.degree. C. for 30 minutes, suggesting that the protective substances are protein in nature. Since a significant increase in anti-I. multifiliis antibody activity was found in plasma, mucus and tank water after the secondary immunization of the hardy fish the protective substances in the water must be the antibodies released from the immunized fish through mucus. Although skin mucus was able to kill certain parasites was reported by Ingram, G. A., "Substances involved in the natural resistance of fish to infection--a review", J. Fish Biol., 16, (1980) 23-60 and specific immunoglobulins were found in the mucus layer of some fish (Di Conza J. J. et al., "Relationship of catfish serum antibodies to immunoglobulin in mucus secretions", Aust. J. Exp. Biol. Med. Sci., 49, (1971), 517-519; Ourth, D. D., "Secretory IgM, Lysozyme and lymphocytes in the skin mucus of the channel catfish, Ictalurus Punctatus", Develop. Comp. Immunol., 4, (1980) 65-74) we are the first group to demonstrate that skin mucus of immunized fish could release antibodies into the water thereby producing effective protection against pathogens concerned for naive or infected fish.

TABLE 3 Anti-I.
multifiliis antibody titres in serum, mucus and tank water of double-deck aquaria containing a catfish pre-injected with 1 or 2 doses of I. multifiliis. Group (catfish in Antibody titres (-log.sub.2) lower tank) n Serum Mucus Tank Water

							Control	9	2.05	+-	0.43
0.00	+-	0.00	0.00	+-	0.00	(naive fish)	Test	9	2.52	+-	0.87
0.00	+-	0.00	0.00	+-	0.00	(1 wk after 1st injection)	Test	9			
5.45	+-	0.65	sup.a	1.65	+-	0.82	sup.d	0.00	+-	0.00	(3 wk after 1st injection)
						Test	9	7.22	+-	0.45	sup.b
0.34	sup.e	0.00	+-	0.00		(5 wk after 1st injection)	Test	9	9.16		
+-	0.45	sup.c	4.05	+-	0.46	sup.e	0.13	+-	0.76	sup.g	(1 wk after 2nd injection)
						Test	9	11.87	+-	0.35	sup.c
0.72	sup.f	0.95	+-	0.08	sup.h	(2 wk after 2nd injection)					

n = number of replicates.
.sup.a,d,g p < 0.05, .sup.b,e,h p < 0.005 and .sup.c,f p < 0.001 significantly different from their controls. Different superscripts indicate significant difference (p < 0.05) between test groups.

.sup.a,d p < 0.05,
 .sup.b,e p < 0.005 and .sup.c p < 0.001 significantly different
 from their controls. Different superscripts indicate significant
 difference (p < 0.05) between test groups.

TABLE 6

The, in vitro,
 immobilization of aquarium tank water on active tomites of I.
 multifiliis as in Table 5 except the lower tank containing 2
 shubankin which were pre-injected with 1 or 2 doses of I.
 multifiliis. Group No. of immobilized tomites (%) (shubankin in
 Days lower tank) 4 8 12

Control	0.0	+-.	0.0	0.0	+-.	0.0	0.0	+-.	0.0	(naive fish) Test
2.2	+-.	1.4	3.3	+-.	1.7	5.6	+-.	1.8	(1 wk after 1st injection)	
Test 6.6	+-.	1.7	.sup.a	16.7	+-.	2.9	.sup.b	27.8	+-.	2.8
(3 wk after 1st injection)										
2.0	.sup.b	36.7	+-.	2.3	.sup.c	(5 wk after 1st injection)				
Test 30.0	+-.	3.7	.sup.c	36.7	+-.	3.3	.sup.c	46.7	+-.	3.3
after 2nd injection)										
Test 55.5	+-.	1.7	.sup.e	(2 wk after 2nd injection)						

.sup.a,d p < 0.05,
 .sup.b,e p < 0.005 and .sup.c p < 0.001 significantly different
 from their controls. Different superscripts indicate significant
 difference (p < 0.05) between test groups.

The above findings clearly demonstrate that antibodies are
 released into the aquatic environment by the immunized fish. It
 is without doubt that these substances not only kill the
 parasites or neutralize their toxins (if any) in the tank water
 but also removing parasites from the fish. The decrease of the
 parasites in the aquatic environment and on the fish body surface
 might allow the infected fish to have sufficient time to build up
 their active immunity.

EXAMPLE 3

Tables 8 and 9 show the effectiveness of the closed aquarium
 system (FIG. 2) when there were more tanks and more infected fish
 of different species. Pre-immunized fish may be in one or more
 tanks. The data again clearly demonstrate that a significantly
 greater number of infected fish recovered from infection with
 immunized fish in a separate tank as compared to the controls,
 irrespective of species. Hence, the present method must also work
 for various species of larger fish which were demonstrated to be
 infected by similar pathogenic protozoans (Hines, R. S. et al.,
 "Ichthyophthiriasis in the mirror carp *Cyprinus carpio*", III.
 Pathology, J. Fish Dis., 6, (1973) 189-196; Goven, B. A., et al.,
 "Protection of channel catfish, *Ictalurus punctatus* Rafinesque,
 against *Ichthyophthirius multifiliis* Fouquet by immunization", J.
 Fish Biol., 17, (1980) 311-316).

In the method of the present invention the pre-immunized fish and
 the naive or infected aquatic lifeforms may be in a flow-through
 method, where the number ratio of pre-immunized fish to fish to
 be protected is increased to a level compatible with producing
 sufficient substances effective against the specific pathogenic
 organisms. The pre-immunized fish and aquatic lifeforms to be
 protected may be, alternatively, in a floating net-cage. The

Alternatively, the fish to be pre-immunized are species which tolerate injections of all kinds of vaccines and are able to respond by producing substantial amounts of substances effective against the pathogens concerned. The method of the present invention is applicable in hatcheries wherein the lifeforms to be protected are at a developmental stage that have an immature immune system.

TABLE 7

multifiliis antibody activity of aquarium water containing naive or immunized fish after heat treatment. The immunized fish were pre-injected with 1 or 2 doses of I. multifiliis. Immobilization test of Group undiluted aquarium water* (catfish in No heat 47.degree. C. 100.degree. C. lower tank) n treatment 30 min. 30 min.

Anti-I.	
(naive fish) Test 12 - - - (1 wk after 1st injection)	Control 12 - - -
- (3 wk after 1st injection) Test 12 + + - (5 wk after 1st injection)	Test 12 + + -
Test 12 + + - (1 wk after 2nd injection)	Test 12 + + -
(2 wk after 2nd injection)	

n = number of replicates. +: Tomites immobilized. -: Motility of tomites not affected. *: Determined on 7th day after fish being put into the closed aquarium system.

TABLE 9

Control of Ich on infected fish, 10 in each upper tank, by 2 immunized shubankin in the lower tank of the closed aquarium system (FIG. 2). The immunized shubankin were obtained at 2 wk after the second injection of the live tomites of *I. multifiliis*. Survival rate of infected fish (%) Lower tank containing Infected immunized Fish n

No fish	Naive Fish	Fish fish**										
Angelfish	3	33.3	30.0	83.3	Goldfish	3	30.0	26.6	86.7	Platy	3	20.0
23.3	80.0	Tiger barbs	3	3.3	6.7	56.7						

n = number of replicates (total 90 fish). **significantly ($p < 0.005$) different from their controls.

4/2/01 7:38 AM

The next two experiments were to test the method with ornamental fishes collected from two fish farms where the fishes were found to be carriers of some infectious diseases. Different numbers of the 3 species were kept in the closed aquarium system. The results in Tables 10 and 11 showed that significant protection was found when they were kept with the immunized fish. Fish in the controls died of white-spot (Ich) or velvet disease which was due to the spread of those protozoans from carrier fish. The ornamental fishes which were kept in the closed culture system with immunized fish remained healthy and free of parasitic protozoan diseases for a period of 6 months before the experiment was terminated.

This shows that the method does not provide a transient protection but that it lasts as long as the particular pathogens are still present in the water. The study also showed that fish from the above fish farm kept in the closed culture system with a secondary immunized catfish against *I. multifiliis* had a significantly higher survival rate towards other protozoan parasites (*Ichthyopodo*, *Trichodina* and *Chilodonella*) as compared to the controls (Table 12). Our studies also showed that fish recovered from "ich" exhibited effective protection against velvet disease. In addition, goldfish immunized with *I. multifiliis* developed protective immunity not only against *I. Multifiliis* but also towards other ectoparasitic protozoans, such as *Ichthyobodo necatrix*, *Chilodonella cyprini*, and *Trichodina* sp., which are commonly found in the tropics (Tables 13-17). The results also showed that the protection was correlated with the antibody titers in their plasma and skin mucus (Table 18).

TABLE 10 Control of Ich and velvet disease on ornamental fish collected from fish farm A*
in a closed culture system using a secondary immunized catfish. Each of the 3 upper tanks consisted of different number of different species of ornamental fish. The lower tank contained a catfish which was pre-injected with 2 doses of *I. multifiliis*. Survival Rate (%) No. of fish of each species used Control.

Test	sup.1	Fish**	10	20	30	10	20	30	
Goldfish	20	15	23.3	100	90				
90 Tiger barbs	10	10	6.7	90	85	86.7	Platy	30	30
								33.3	100
								95	93.5

*Some fish were randomly chosen for parasite determination before putting them into the closed system: Tiger barbs were the carrier of *I. multifiliis* and *Oodinium pillularis* while goldfish were carrier of *Costia* sp. and *Gyrodactylus* sp. **Each species placed separately in a tank.
.sup.1 Significantly ($p < 0.05$) different from their control groups.

TABLE 11 Control of Ich on ornamental fish collected from fish farm B* in a closed culture system using a secondary immunized catfish against *I. multifiliis*. Survival Rate (%) No. of fish of each species used Control Test.

sup.1	Fish**	10	20	30	10	20	30	
Goldfish	20	30	30	90	100			
92.3 Tiger barbs	30	25	26.3	100	90	96.6	Platy	20
								10
								90
								85
								86.6

*Platy were the carrier of

I. multifiliis and Tricodina sp. Goldfish were the carrier of Costia sp. and Tricodina sp. Tiger barbs were the carrier of Costia sp. and Gyrodactylus sp. **Each species placed separately in a tank. .sup.1 Significantly ($p < 0.05$) different from their control groups.

TABLE 12 Control of
Chilodonella, Ichthyopodo, and Trichodina on ornamental fish
collected from fish farm B in a closed culture system using a
secondary immunized catfish against I. multifiliis. Survival Rate
(%) Chilodonella Ichthyopodo Trichodina Fish** Control Test
Control Test Control Test

Goldfish	6.7	93.3	24.5	85.6	13.3	87.5	Tiger barbs	3.3	87.5	6.3
	76.5	15.5	82.6	Platy	10	95.5	15.5	90.5	25.5	92.6

n = 3 replicates (total 30 fish). **Each species placed separately in a tank. .sup.1
Significantly ($p < 0.05$) different from their control groups.

EXAMPLE 5

Table 19 shows that immunized hardy fish when kept with the bacteria-infected goldfish showing fin- and gill-rot disease were able to facilitate the recovery of the latter. On the contrary, all the infected goldfish when kept with the naive hardy fish died within 14 days.

EXAMPLE 6

Tables 20 and 21 show the effectiveness of the closed aquarium system (FIG. 2) when there were more tanks and fish fry of different ornamental fish species were kept. The data again clearly demonstrate that a significantly higher protection on fry kept with the immunized catfish in a separate tank as compared to the controls, irrespective of species. Hence, the present method also works for younger stages (fry) of different ornamental fish species.

The above detailed examples (1 to 6) of the present invention demonstrate that the method provides a relatively pathogen-free closed aquatic environment for naive or infected ornamental fish in which pathogens are kept in check by antibodies released from hardy fish which have been immunized against those pathogens. This method is self-generating with antibodies continuously and increasingly being released by the immunized fish as long as the particular pathogens are still present in the water (a process known as secondary immune response). By this means, the antibodies constantly act as a preventive measure for the aquatic environment by removing, or suppressing the population of, pathogens in the water, thereby preventing infection or further infection or disease outbreak.

Furthermore, a variety of antibodies against various common infectious pathogens in a particular species of aquatic lifeform can be introduced into the aquatic environment at the same time by immunizing a small number of hardy fish with those vaccines. For instance, a few hardy fish can be immunized with a few vaccines of pathogenic protozoans, bacteria and viruses that

commonly infect culture fish. Thus, antibodies against these pathogens will be released into the aquatic environment. This method is more effective and practical than by injecting heterologous antibodies into fish for passive immunity (Frechette et al., U.S. Pat. No. 5,064,645, issued Nov. 12, 1991). The latter method would require repeated administration whereas the present invention is a self-generating system with antibodies continuously and increasingly being released by the immunized fish as long as the particular pathogens are still present in the water.

TABLE 13 Protection of
naive and immunized goldfish against *I. multifiliis* challenge. The immunized fish had been treated twice with live tomites of *I. multifiliis* by immersion or injection. No. not infected Group
Total no. Protection Mortality (treatment) challenged (%) (%)

(naive goldfish) Test 1** 90/90 100.0 ^{sup.a} 0.0 ^{sup.b} (immersion) Control 2* 3/90 3.3 95.6 naive goldfish Test 2** 88/90 97.8 ^{sup.a} 0.0 ^{sup.b} (injection)	Control 1* 2/90 2.2 97.7
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------

*Naive (control) goldfish had been injected with or immersed twice in saline prior to challenge. **A very small number of these fish showed light infections. ^{sup.a}Percentage of protection is significantly ($P < 0.05$) different from controls. ^{sup.b}Percentage of mortality is significantly ($P < 0.05$) different from controls.

TABLE 14 Protection of
naive and immunized goldfish against *O. pillularis* challenge. The immunized fish had been treated twice with live tomites of *I. multifiliis* by immersion or injection. No. not infected Group
Total No. Protection Mortality (treatment) challenged (%) (%)

(naive goldfish) Test 1** 70/90 77.7 ^{sup.a} 11.1 ^{sup.b} (immersion) Control 2* 1/90 1.1 100.0 naive goldfish Test 2** 66/90 73.3 ^{sup.a} 13.3 ^{sup.b} (injection)	Control 1* 0/90 0.0 100.0
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------

*Naive (control) goldfish had been injected with or immersed twice in saline prior to challenge. **A small number of these fish showed light infections. ^{sup.a}Percentage of protection is significantly ($P < 0.05$) different from their controls. ^{sup.b}Percentage of mortalities is significantly ($P < 0.05$) different from their controls.

TABLE 15 Protection of
naive and immunized goldfish against *Trichodina* challenge. The immunized fish had been treated twice with live tomites of *I. multifiliis* by immersion or injection. No. not infected Group
Total No. Protection Mortality (treatment) challenged (%) (%)

naive goldfish Test 1** 46/60 76.7 ^{sup.a} 8.3 ^{sup.b} (immersion) Control 2* 2/60 3.3 86.7 naive goldfish Test 2** 42/60 70.0 ^{sup.a} 13.3 ^{sup.b} (injection)	Control 1* 0/60 0.0 83.3
--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------

*Naive (control) goldfish had been injected with or immersed twice in saline prior to challenge. **A small number of these fish showed light infections. ^{sup.a}Percentage of protection is significantly ($P < 0.05$) different from controls. ^{sup.b}

Percentage of mortality is significantly ($P < 0.05$) different from controls.

TABLE 16

Protection of naive and immunized goldfish against Ichthyobodo challenge. The immunized fish had been treated twice with live tomites of I. multifiliis by immersion or injection. No. not infected Group Total No. Protection Mortality (treatment) challenged (%) (%)

Group	Test	1**	45/60	75.0	sup.a	8.3	sup.b	(immersion)
naive goldfish	Control 1*	0/60	0.0	80.0				
naive goldfish	Control 2*	1/60	1.7	75.0				
naive goldfish	Test 2 a**	44/60	73.3	sup.a	5.0	sup.b		(injection)

*Naive (control) goldfish had been injected with or immersed twice in saline prior to challenge. **A small number of these fish showed light infections. .sup.a Percentage of protection is significantly ($P < 0.05$) different from controls. .sup.b Percentage of mortality is significantly ($P < 0.05$) different from controls.

TABLE 17

Protection of naive and immunized goldfish against Chilodonella challenge. The immunized fish had been treated twice with live tomites of I. multifiliis by immersion or injection. No. not infected Group Total No. Protection Mortality (treatment) challenged (%) (%)

Group	Test	1**	28/30	93.3	sup.a	0.0	sup.b	(immersion)
naive goldfish	Control 1*	3/30	10.0	66.7				
naive goldfish	Control 2*	5/30	16.7	83.3				
naive goldfish	Test 2**	27/30	90.0	sup.a	3.3	sup.b		(injection)

*Naive (control) goldfish had been injected with or immersed twice in saline prior to challenge. **A small number of these fish showed light infections. .sup.a Percentage of protection is significantly ($P < 0.05$) different from controls. .sup.b Percentage of mortality is significantly ($P < 0.05$) different from controls.

TABLE 18

Serum and mucus antibody titres of naive and immunized goldfish which were Immunized twice by live tomites of I. multifiliis (immersion or injection) against different parasitic protozoans. Antibody titres (-log.sub.2) Test Control (immunized Group (Naive Fish) fish) (treatment) Mucus Plasma Mucus Plasma

Group	Test	1**	5.98	sup.a	10.65	sup.a	2.00	sup.a	4.43	sup.a	11.11	sup.a	0.00	0.44	sup.e	0.59	sup.a	0.00	0.22	sup.e	3.33	sup.e	5.62	sup.e	0.00	0.67	sup.e	0.79	sup.c	0.00	0.58	sup.e	0.39	0.86	sup.e	0.43	sup.c	0.11	sup.e	0.89	sup.e	0.51	0.69	sup.e	0.67	sup.c	0.00	0.36	sup.f	1.15	sup.c	0.00	0.33	sup.f	2.22	sup.f	5.01	sup.f	0.00	0.52	sup.f	0.43	sup.c	0.11	sup.e	0.81	sup.b	0.00	1.79	sup.b	3.52
naive goldfish	Anti-Ich.*	0.00	5.98	sup.a	10.65	sup.a	2.00	sup.a	4.43	sup.a	11.11	sup.a	0.00	0.44	sup.e	0.59	sup.a	0.00	0.22	sup.e	3.33	sup.e	5.62	sup.e	0.00	0.67	sup.e	0.79	sup.c	0.00	0.58	sup.e	0.39	0.86	sup.e	0.43	sup.c	0.11	sup.e	0.89	sup.e	0.51	0.69	sup.e	0.67	sup.c	0.00	0.36	sup.f	1.15	sup.c	0.00	0.33	sup.f	2.22	sup.f	5.01	sup.f	0.00	0.52	sup.f	0.43	sup.c	0.11	sup.e	0.81	sup.b	0.00	1.79	sup.b	3.52

+- . 8.56 +- . (injection) 0.00 0.51 0.33.sup.e 0.88.sup.b
 *: Immobilization test;
 **: Inhibition of division. .sup.a,d : P < 0.005, .sup.b,e p < 0.01 and .sup.e,f p < 0.05: significantly different from controls. Different superscripts indicate significantly (p < 0.05) different between test groups.

TABLE 19 Control of fin rot caused by Aeromonas hydrophila in goldfish by catfish or shubankin immunized with formalin-killed A. hydrophila on day 0 and 14. Two weeks after the 2nd injection, immunized fish were placed in the closed culture system. No. of Group recovered Survival (fish in lower No. of rate tank) n infected (%)
 Control 1 3 0/30 0 (no fish) Control 2 3 0/30 0 (naive catfish) Test 1 3 25/30 83.3.sup.a (immunized catfish) Test 2 3 22/30 73.3.sup.b (immunized shubankin) n = number of replicates. .sup.a,b p < 0.005 significantly different from the controls.

TABLE 20 Control of Ichthyophthiriasis in goldfish fry in a closed recirculation system by using secondary immunized catfish. No. not No. of Average infected recovered Survival Total Group Total no. No. of rate Protection (treatment) challenged infected (%) (%)
 I 0/270 18/270 6.7 6.7
 Control 1* (no fish in lower tank) II 3/270 22/267 8.2 9.3
 Control 2* (naive catfish in lower tank) III** 118/270 96/152 63.2.sup.a 79.3.sup.b (immunized catfish in lower tank)
 *Inflicted of heavy infection of I. multifiliis. **About half of the naive goldfish fry showed light infection. .sup.a Percentage of survival is significantly (P < 0.005) different from the controls. .sup.b Percentage of total protection is significantly (P < 0.005) higher from the controls.

TABLE 21 Control of Ichthyophthiriasis in guppy and platy fry in a closed recirculation system by using secondary immunized catfish. No. not No. infected recovered Average Total Group Total no. No. Survival Protection (treatment) challenged infected (%) (%)
 I (guppy) 2/180 33/178 18.5 19.4 Control 1 (no fish in lower tank) II (guppy)* 3/180 35/267 19.4 21.1 Control 2 (naive catfish in lower tank) III (guppy)** 96/180 63/84 75.0.sup.a 88.3.sup.b (immunized catfish in lower tank) IV (platy)* 0/90 8/90 8.9 8.9 Control 1 (no fish in lower tank) V (platy)* 0/90 10/90 11.1 11.1 Control 2 (naive catfish in lower tank) VI (platy)** 42/90 29/48 60.4.sup.a 78.9.sup.b (immunized catfish in lower tank)
 *Inflicted of heavy infection of I. multifiliis. **About half of the naive fish showed light infection. .sup.a Percentage of survival is significantly (P < 0.005) different from the controls. .sup.b Percentage of total protection is significantly (P < 0.005) higher from the controls.

Further Applications of the Inventive Concept

Although we have tested the invention for the control of some protozoan diseases and one bacterial disease in freshwater fishes, the inventive concept can be applied for the control of any disease (whether due to parasites, bacteria, viruses or fungi) in any aquatic species (whether freshwater or marine and finfish or shellfish such as crustaceans and mollusks, or other lifeforms). The method to be applied to any aquatic species for the control of any disease will be the same as described above. Let us take the example of species A, which may be any species of crustacean, mollusk, fish, or any other aquatic lifeform (e.g. sea urchins), whether freshwater or marine, and whether eggs, larvae, fry, juveniles or adults. If we know the pathogens causing diseases in species A and can obtain the corresponding vaccines or the isolates to prepare the vaccines by available methods, the vaccines can be injected into any species of fish known to be hardy and to show good immune response, preferably with a good capacity to produce mucus that live in the same type of environment (i.e. freshwater or marine) as species A. At least two injections, spaced 14 to 30 days apart, are recommended to ensure a good secondary immune response. These fish will develop antibodies against the pathogens concerned (even if these pathogens of species A are not normally pathogenic to fish. Anyhow, the fish will produce an immune response as long as these pathogens are antigens to the fish).

These immunized fish are then kept with species A either in the same enclosure or in a separate enclosure but connected with the enclosure of species A in terms of the water flow. The immunized fish release antibodies into the water, and as long as the immunized fish continue to encounter the pathogens concerned in the water they will develop and release an increasing amount of antibodies into the water, thereby preventing infection/further infection and outbreak of the disease(s) in species A, and even causing rapid recovery in diseased individuals of species A.

The general principle is that any organism which causes a disease in any aquatic species of any stage of development, may be prepared as a vaccine for injection into hardy fish selected for this purpose even though the organism may not be pathogenic in the fish itself. As a foreign body, the organism-derived vaccine will elicit an immune response in fish thereby causing the production of antibodies which are released into aquatic environment continuously and in increasing quantities as long as the pathogen is still present in the water. These antibodies in the water will protect against infection and disease outbreak in that particular species.

It is to be noted that the inventive method is to make use of the immunized hardy fish, and not the cultured shellfish or other aquatic species, to produce and release protective antibodies, into the aquatic environment, to kill the particular pathogens in the water or those infecting the cultured lifeforms. Therefore, the inventive method can be used to control diseases in not only fish but also other aquatic lifeforms regardless of the species, size or developmental stage. The immunized fish are chosen as the source of antibodies because fish have been shown to produce

protective humoral and cell-mediated immune response towards various aquatic disease-causing organisms such as *Aeromonas hydrophila*, *Edwardsiella ictaluri*, Infectious pancreatic necrosis, *Vibrio* and many others (Ellis, A. E., Fish Vaccination. Academic Press, London (1988)). Even if the pathogens that infect crustaceans, mollusks, echinoderms and other aquatic lifeforms are antigenically different from fish pathogens, the immune system of fish can still respond to the pathogens, and therefore the immunized fish can still be used as the source of antibodies for release into the water for protecting the said aquatic lifeform against the pathogens. At the minimum, the inventive concept can be used to effectively reduce or eliminate pathogenic organisms in the aquatic medium through placing in the medium appropriate fish which have been immunized against those pathogenic organisms.

In conclusion, the results of our experiments demonstrate that immunized hardy fish such as catfish and shubankin in a closed system of culture with infected fish are effectively enhancing the recovery of the latter, irrespective of species. The data also strongly evidence that the protection is due to a high level of anti-I. multifiliis antibodies present in the mucus of immunized fish which is released into the circulating water. The anti-I. multifiliis antibody activity was shown to protect fish not only from "Ich" but also against Oodinium and other protozoans, such as Ichthyobodo, Trichodina and Chilodonella. Since the 3 species of ornamental fish kept in this system with a immunized catfish were free of infectious protozoan diseases for a period of 6 months before the experiments were terminated despite some fish being carriers of infectious protozoa (Tables 10 and 11), the method is also effective in controlling the spread of the infectious organisms as long as needed. The method works equally well with a bacterial disease as shown in Table 19 and also prove to be very effective for the protection of fry of different species of ornamental fish against infectious parasitic diseases (Tables 8 and 9). All these data so far obtained have demonstrated that the immunized fish in a closed culture system is highly effective in protecting naive fish against infectious disease-causing organisms as well as facilitating the recovery of infected fish. Hence, it should work for any infectious disease, be it parasitic, bacterial, viral or fungal.

Although the efficacy of the present inventive method has been repeatedly tested only in ornamental fish, the inventive concept is applicable to other fish species as well as crustaceans, mollusks and other aquatic lifeforms (e.g. echinoderms), whether freshwater or marine and irrespective of the developmental stage and size. This is based on the following facts:

- (1) Fish can produce protective substances (antibodies) against various vaccines prepared from pathogenic organisms, even those which are not pathogenic in the fish themselves (prior art, see references)
- (2) The immunized fish can release the protective substances into the water (present invention).

(3) The immunized fish will continuously and increasingly release protective substances into the water as long as the pathogenic organisms are still encountered (present invention),

(4) The released protective substances can kill the pathogenic organisms in the water, thus removing them from the water (present invention).

(5) The released protective substances can even effect recovery of infected individuals (present invention).

It will be understood that the present invention is not limited to the method described but includes all modifications and variations that may come within the scope of the appended claims.

CLAIMS:

We claim:

1. A method of controlling one or more pathogenic organisms selected from the group consisting of protozoan parasites and bacteria wherein the pathogenic organisms are detrimental to aquatic lifeforms selected from the group consisting of fish, crustaceans, mollusks and echinoderms wherein the method comprises;

a) adding immunized fish immunized against one or more of said pathogenic organisms to a culture system comprising an aquatic medium;

b) permitting said immunized fish to release antibodies against the pathogenic organisms into the aquatic medium;

c) providing the aquatic lifeforms to the aquatic medium wherein the aquatic lifeforms are to be treated or protected; and

d) allowing the antibodies to immunoreact with the pathogenic organisms, thereby controlling the pathogenic organisms.

2. The method of claim 1, wherein the immunized fish have been injected at least once or twice with pathogenic organisms selected from the group consisting of Ichthyophthirius, causing white-spot disease, Oodinium, causing velvet disease, Ichthyobodo, Trichodina and Chilodonella, causing slimy skin, and Aeromonas, causing fin-rot and gill-rot disease.

3. The method of claim 1, wherein the culture system comprising a closed recirculation culture system incorporating a tank containing the immunized fish.

4. The method of claim 1, wherein the immunized fish are in one tank and the aquatic lifeforms are contained in one or more tanks connected to the tank containing the immunized fish and wherein the aquatic medium in the tank containing the immunized fish is recirculated between the one or more tanks containing the aquatic lifeforms.

5. The method of claim 1, wherein the immunized fish are in a flow-through tank with the aquatic lifeforms, and wherein the number ratio of immunized fish to aquatic lifeforms to be protected is sufficient to provide an effective amount of antibodies protective against the pathogenic organisms.
6. The method of claim 1, wherein the immunized fish and aquatic lifeforms are in a floating net-cage.
7. The method of claim 1, wherein the immunized fish are species which tolerate injections and are capable of producing body mucus containing antibodies which are effective against pathogenic organisms.
8. The method of claim 1, wherein the immunized fish and the aquatic lifeforms are both freshwater organisms.
9. The method of claim 1, wherein the immunized fish and aquatic lifeforms are both marine organisms.
10. The method of claim 1, wherein the aquatic lifeforms are at any stage of development.
11. The method of claim 8, wherein the aquatic lifeforms to be protected are non-infected.
12. The method of claim 1, wherein the aquatic medium is a component of a quarantine system for treating the aquatic lifeforms before culture or export.
13. The method of claim 10, wherein the aquatic medium is a component of a hatchery.
14. The method of claim 1, wherein the immunized fish are selected from the group consisting of catfish and shubankin.
15. The method of claim 14, wherein the aquatic lifeforms are selected from the group consisting of freshwater angel fish, goldfish, platy, and tiger barb.
16. The method of claim 15, wherein pathogenic organisms are selected from the group consisting of protozoans and bacteria wherein the protozoans cause white-spot disease, velvet disease and slimy skin disease, and wherein the bacteria cause fin-rot or gill-rot disease.
17. The method of claim 1, wherein the culture system is one or more selected from the group consisting of a closed recirculation system, a flow-through system, a floating net-cage, a pond system, a hatchery, a grow-out system, a tank, a holding tank and a quarantine tank.
18. The method of claim 3, wherein the immunized fish and the aquatic lifeforms are in the same tank in the closed recirculating culture system.